



SC-07

Role of health education on HPV vaccination willingness among a sample of students in the Local Health Authorities of Agrigento, Palermo and Trapani, Sicily

F. Cacioppo (1), C. Amodeo (1), N. Bonaccorso (1), M. Sciortino (1), E. Zarcone (1), M. Tocco (1), L. Pipitone (1), A. Carubia (1), L. D'Azzo (1), T. Vitello (1), G. Bosco (2), S. Palmeri (3), V. Di Gaetano (4), G. Canzoneri (4), F. Di Gregorio (4), L. Vitale (4), L. Vinci (4), A. Graceffa (2), V. Spoto (2), D. Mirabile (3), R. Asciutto (3), S. Paolizzo (3), W. Mazzucco (1), F. Vitale (1), A. Casuccio (1), E. Amodio (1), A. Pieri (1), G. Graziano (1), C. Costantino (1)

(1) Department of Health Promotion Sciences, Maternal and Infant Care, Internal Medicine and Excellence Specialties "G. D'Alessandro", University of Palermo; (2) Department of Prevention, Local Health Authority of Agrigento; (3) Department of Prevention, Local Health Authority of Palermo; (4) Department of Prevention, Local Health Authority of Trapani.

Background. HPV is one of the most common causes of sexually transmitted infections (STIs) worldwide. Strategies to counteract HPV-related infections and diseases include primary prevention measures, such as vaccination, and secondary prevention measures, such as Pap test and HPV DNA test. The objective of this study is to assess the extent to which health education and knowledge levels influence students' propensity to make informed health choices, with particular reference to HPV vaccination.

Materials and Methods. A total of 2,111 middle and high school students from the provinces of Agrigento (1 schools), Palermo (2 schools), and Trapani (26 schools) participated in the study. They were administered a questionnaire designed to assess their knowledge of STIs, particularly HPV, as well as their attitudes toward and adherence to HPV vaccination. Participation in the study was contingent upon the parents receiving the information sheet and signing the informed consent form.

Results. The sample consisted of 780 (36.3%) males and 1,371 (63.7%) females. 64.6% of participants reported being vaccinated against HPV, 22.6% reported not being vaccinated, and 12.8% reported not knowing their vaccination status; vaccination coverage was similar among males (62.8%) and females (61.6%). The majority of students (79.4%) reported not having yet had sexual intercourse, while 11.8% reported using contraceptive methods. 89.4% of students reported having heard of HPV (86.8% of males and 90.8% of females). This awareness is associated with greater willingness to be vaccinated (8.8 vs. 7.3 among those who had never heard of it; p -value <0.001). Similarly, awareness that HPV can cause clinical manifestations is associated with higher levels of willingness: 9.02 among aware students, 8.11 among those who were unsure, and 7.8 among those who mistakenly believed that HPV could not cause them (p -value <0.001). Regarding sources of information on STIs, 18.8% of students reported having heard about them at school (willingness 8.76), 38.4% within their families (8.99), and 20.0% through healthcare professionals (9.08). The highest levels of willingness were observed among students exposed to all major sources of information (9.22), while they were lower among those who had not received information in any context (8.12) (p -value <0.001).

Conclusions. The results highlight a clear association between the level of knowledge about STIs and HPV and participation in health education interventions, with a greater propensity for vaccination. In particular, consulting multiple and integrated information sources is associated with the highest levels of willingness to be vaccinated. These data underscore the crucial role of health education in promoting informed health behaviors and in helping to achieve vaccination prevention goals. Strengthening educational programs, especially in schools, can help improve knowledge of HPV and related diseases, with a positive impact on HPV vaccination uptake.